

## Well Child Check-Up and Immunization Schedule (Birth-S years)

Each check-up includes a growth and weight evaluation, each well child appointment includes a full developmental assessment and a complete physical examination.

### Age at Check-Up:

- 3 to 5 days
- 3 Weeks
- 6 Weeks\*
- 4 Months
- 6 Months
- 9 Months
- 12 Months\*
- 15 Months\*
- 18 Months
- 2 Years
- 2 Years
- 3 years
- 4 Years\*
- 5 years

### Immunizations/Screening Tests:

- Post Hospital Exam and Weight Check follow ups as indicated
- Physical Exam and Weight Check follow up
- DTap,HepB,Ipv {Pedarix},Hib, PCV13 (prevnar), RV1/5 (Rotavirus)
- DTap,Hib,Ipv {Pentacel}, PCV13 (prevnar), RV1/5 (Rotavirus)
- DTap,HepB,Ipv {Pedarix},Hib, PCV13 (prevnar), RV1/5 (Rotavirus)
- Sickle Cell Screen/Lead Screen/HGB-HCT Screen/Developmental Screen
- DTap,Hib,Ipv (Pentacel), PCV13 (prevnar), MMR,VAR, HepA, TB Screen
- HGB-HCT Screen/Lead Screen
- Physical Exam and Weight Check follow up
- Physical Exam and Weight Check follow up, HepA, Developmental Screen
- Physical Examination and Weight Check Follow up
- Physical Examination and Weight Check follow up, HGB-HCT Screen
- Lead screen
- Physical Examination, Vision Screen, Hearing Screen, Urinalysis
- DTap,Ipv (Kinrix), MMRV (proquad), Vision Screen, Hearing Screen
- TB Screen, Physical Examination, and Urinalyiss
- Physical Exam, Hearing Screen, Vison Screen and Urinalysis

\*\* The 6 weeks, 12 months, 15months, and 4 years exams are to be done on time. After 5 year, well child checks are recommended every year. Universal Pediatrics does require yearly check-ups/physicals. For patients with chronic problems, such as Asthma and ADHD we require that the patients are seen every 3 months.

### Abbreviations Used:

- Dtap: Diphtheria, Tetanus, Acellular Pertusis (Whooping Cough)

- Tdap: Tetanus, Diphtheria, Acellular Pertusis
- Hep B
- Hep A
- Hib
- Ipv
- MMR
- Var
- TB
- Prevnar
- Rotavirus
- Hepatitis B Vaccine
- Hepatitis A Vaccine
- Haemophilus Influenzae, type B (bacteria that can cause meningitis)
- Inactivated Polio Vaccine
- Measles, Mumps, Rubella
- Varicella (Chicken Pox)
- Tuberculosis
- Pneumococcal Vaccine (bacteria that can cause meningitis)
- Oral Vaccine against a common intestinal virus
- MCV4 Meningitis
- HPV Human Papilloma Virus Vaccine

**Adolescent Vaccines:**

- MCV is given between 11-16 years old, one booster after 16 years old before college entry
- Tdap is given between 11-13 year old
- Hpv is given in 3 doses starting at 9 years old for both male and female patients